

Magnetic Pigment

INSAT
Ins. 1
Ins. 2

Subject matter of the invention are magnetic particles having a glass surface, and a procedure for purifying a biological material, especially nucleic acids, using glass particles in the presence of chaotropic salts. Yet another subject matter of the invention is a procedure for isolating these biological materials and a procedure for concentrating biological materials and transferring them from solutions having a high concentration of salts to solutions having a low concentration of salts.

Add B3

Many biological materials, especially nucleic acids, present special challenges in terms of isolating them from their natural environment. On the one hand they are often present in very small concentrations and, on the other, they are often found in the presence of many other solid and dissolved substances that make them difficult to isolate or measure.

For this reason, many procedures and materials for isolating nucleic acids from their natural environment have been proposed in recent years. In Proc. Natl. Acad. USA 76, 615-691 (1979), for instance, a procedure for binding nucleic acids in agarose gels in the presence of sodium iodide in ground flint glass is proposed.

The purification of plasmid DNA from bacteria on glass dust in the presence of sodium perchlorate is described in Anal. Biochem. 121, 382-387 (1982).

In DE-A 37 34 442, the isolation of single-stranded M13 phage DNA on glass fiber filters by precipitating phage particles using acetic acid and lysis of the phage particles with perchlorate is described. The nucleic acids bound to the glass fiber filters are washed and then eluted with a menthol-containing buffer in Tris/EDTA buffer.

A similar procedure for purifying DNA from lambda phages is described in Anal. Biochem. 175, 196-201 (1988).

The procedure known from the prior art entails the selective binding of nucleic acids to glass surfaces in chaotropic salt solutions and separating the nucleic acids from contaminants such as agarose, proteins or cell residue. To separate the glass particles from the contaminants according to the prior art, the particles are either centrifuged or fluids are drawn through glass fiber filters. This is a limiting step, however, that prevents the procedure from being used to process large quantities of samples.

The use of magnetic particles to immobilize nucleic acids after precipitation by adding salt and ethanol is described in Anal. Biochem. 201, 166-169 (1992) and PCT GB 91/00212. In this procedure, the nucleic acids are agglutinated along with the magnetic particles. The agglutinate is separated from the original solvent by applying a magnetic field and performing a wash step. After one wash step, the nucleic acids are dissolved in a Tris buffer. This procedure has a disadvantage, however, in that the precipitation is not selective for nucleic acids. Rather, a variety of solid and dissolved substances are agglutinated as well. As a result, this procedure can not be used to remove significant quantities of any inhibitors of specific enzymatic reactions that may be present.

A porous glass in which magnetic particles are embedded is described in US-A-4,233,169.

Magnetic, porous glass is also available on the market that contains magnetic particles in a porous, particular glass matrix and is covered with a layer containing streptavidin. This product can be used to isolate biological materials, e.g., proteins or nucleic acids, if they are modified in a complex preparation step so that they bind covalently to biotin.

add B⁴
The task of the invention was to provide better materials for immobilizing biological materials and a simple procedure for isolating biological materials, especially nucleic acids, that is also suitable for use in routine diagnostic procedures.

Subject matter of the invention are magnetic particles with an outer glass surface that is substantially pore-free, or that has pores with less than 10 nm diameter. Yet another subject matter of the invention are ferromagnetic particles having a glass surface, a

procedure for isolating biological materials, especially nucleic acids, and a procedure for the manufacture of magnetic glass particles.

Add B5
Add E6 Particles, according to the expert, are solid materials having a small diameter. Particles like these are often also referred to as pigments. According of the present invention, those particles are especially suited that have an average particle size of less than 100 μm . More preferably they have an average particle size of between 10 and 60 μm . The distribution of particle size is preferably relatively homogeneous. In particular, there are almost no particles $< 10 \mu\text{m}$ or $> 60 \mu\text{m}$ in size.

Those materials are referred to as magnetic that are drawn to a magnet, i.e., ferromagnetic or superparamagnetic materials, for instance. In addition, those materials that are called softly magnetic are also understood to be magnetic, e.g., ferrites. Especially preferred according to the present invention are ferromagnetic materials, especially if they have not yet been premagnetized. Premagnetization in this context is understood to mean bringing in contact with a magnet, which increases the remanence. Especially preferred are ferromagnetic materials, such as magnetite (Fe_3O_4) or Fe_2O_3 .


An outer surface of a particle is understood to mean the contiguous surface from which perpendicular lines can be drawn outwards towards the particle's environment that do not cut through the particle itself.

A pore is understood to be a recess in the outer surface of the particle. The surface reaches so far into the particle that a perpendicular line drawn in the recess on the surface cuts the particle at least once in the direction of the adjacent environment of the particle. In addition, pores reach into the particle to a depth that is greater than one radius of the pore.

A glass according to the present invention is understood to be an amorphous material that contains silicium. Glass can contain other materials such as

| | |
|--------------------------------|-----------|
| B ₂ O ₃ | (0 - 30%) |
| Al ₂ O ₃ | (0 - 20%) |
| CaO | (0 - 20%) |
| BaO | (0 - 10%) |
| K ₂ O | (0 - 20%) |
| Na ₂ O | (0 - 20%) |
| MgO | (0 - 18%) |
| Pb ₂ O ₃ | (0 - 15%) |

Glass can also contain a smaller percentage (0 - 5%) of a number of other oxides such as Mn₂O₃, TiO₂, As₂O₃, Fe₂O₃, CuO, CoO, etc. Surfaces made of a composition of borosilicate glass, flint glass or silica have proven to be especially effective. Borosilicate glasses, which are especially preferred in terms of nucleic acid yield, have a boroxide content of more than 25%. A glass having a 70/30 composition of SiO₂/B₂O₃ is especially preferred. Especially preferred according to the present invention are glasses that are formed using the gel sol process and then dried and compressed. The basic principles of this process are known and were described, for instance, in C.J. Brinker, G.W. Scherer "Sol Gel Science - The Physics and Chemistry of Sol Gel Processing", Academic Press Inc. 1990, Sol-Gel Optics, Processing and Applications, Lisa C. Klein, Ed., Kluwer Academic Publishers 1994, p. 450 ff., and in DE-A-1941191, DE-A-3719339, DE-A-4117041 and DE-A-4217432. The principle has not been described for magnetic particles to date, however. The fact that the process could be used to create magnetic particles that have very surprising characteristics when used to isolate biological materials, especially nucleic acids, was not expected. In the gel-sol process, alkoxides of network-forming components, e.g., SiO₂, B₂O₃, Al₂O₃, TiO₂, ZrO₂, GeO₂, are combined with oxides and salts of other components, e.g., in an alcohol solution, and then hydrolized. The equation below describes the procedure for making sodium boroaluminium silicate glass:



- 5 -



OH




→

O-
Na⁺

Gel

T



Water is added to begin the hydrolysis process of the starting components. The reaction proceeds relatively quickly because the alkali ions have a catalytic effect on the speed of hydrolysis of the silicic acid ester. Once the gel is formed it can be dried and densified by means of a thermal process to form glass.

The sol : pigment ratio has a considerable effect on the yield of magnetic pigments provided by this invention. The ratio is limited by the fact that the portion of pigment must be so small that the mass created can still be pumped or sprayed. If the portion of the pigment is too small, the fine portion, e.g., of non-magnetic material, becomes too great and causes interference. Ratios of 10 to 25 g pigment : 100 ml sol were found to be useful in terms of pigment yield.

To create a powder, the slurry is preferably sprayed through a nozzle and the aerosol is dried as it falls. The nozzle is preferably heated to speed up the drying of the slurry. Depending on the nozzle geometry, the nozzle temperature is preferably from 120 to 200° C. A compromise is found by utilizing a sufficient evaporation speed but avoiding overheating.

To optimize the yield, the densification temperature should be as high as possible. If it is too high, however, the particles will stick together and form agglomerates that must be

sieved out. Additional treatment of the particles in at too high temperature will result in a loss of magnetic properties. Too high temperatures should therefore be omitted.

A substantially pore-free surface is understood to mean a surface with pores (as described above) covering less than 5%, but preferably less than 2%, and especially preferred, less than 0.1% of its area. If pores are present, they preferably have a diameter of less than 10 nm and, especially preferred, 1 nm.

Especially preferred according to the present invention are particles that contain a mica core coated with TiO_2 and magnetite particles immobilized on it. In this design, the composite material formed is surrounded by the glass layer. Both the core and the magnetite particles are crystalline and non-porous. The spaces on the surface of the mica that are not occupied by the magnetite particles are covered by a glass layer that is thicker than at the tips of the magnetite particles, basically resulting in a non-porous glass surface.

The non-porosity of the magnetic particles is based only on the outer surface and not on the inside of the particle. The particle can therefore be porous on the inside only if the surface is enclosed by a substantially pore-free glass or a glass surface having pores with a diameter of less than 10 nm.

Surprisingly, the magnetic particles provided by the invention are especially suited for isolating biological materials from samples. Long nucleic acids in particular are not destroyed — or only minimally — when they are immobilized on them. In addition, the core material is a natural resource and therefore causes little ecological concern. Moreover, the particles according to the invention are inexpensive and easy to manufacture.

Yet another object of the invention are ferromagnetic particles having a glass surface. Superparamagnetic particles are described in the prior art. It has been demonstrated that ferromagnetic particles covered with a glass surface offer considerable advantages for isolating biological materials. If the ferromagnetic particles have not been brought in contact with a magnetic field, gravity is the only force that can cause them to sediment

out. They can be resuspended easily and quickly by shaking the solution. The sedimentation procedure that does not utilize a magnetic field preferably proceeds more slowly than the immobilization of biological materials on the surface of the particles. This is especially true for nucleic acids. The ferromagnetic particles can be easily collected at a specific location in the sample fluid by means of a magnet. The fluid is then separated from the particles and, therefore, from the immobilized biological materials.

The glass surface of the ferromagnetic particles provided by the invention can be pore-free or contain pores. For the reasons given above for the magnetic particles provided by the invention, it is preferable for the outer surface of the ferromagnetic particles to also be substantially pore-free or to have pores with a diameter of less than 10 nm. The ferromagnetic particles provided by the invention also preferably have a particle size of between 10 and 60 μm , and especially preferred, of between 20 and 50 μm . Especially preferred are particles with surface pores (if present) having a diameter of less than 10 nm and, especially preferred, 1 nm. An example of a ferromagnetic particle according to the invention is the composite material described above which is made of mica and magnetite particles surrounded by a glass layer.

Yet another object of the invention is a procedure for isolating a biological material by

- bringing a sample containing the biological material in a fluid in contact with the magnetic particles according to the invention or the ferromagnetic particles according to the invention under conditions in which the biological material binds to the particle surface, and
- separating the biological material from the fluid.

Biological materials are understood to mean materials with a particular or molecular basis. They include, in particular, cells such as viruses or bacteria, as well as isolated human and animal cells such as leucocytes, and immunologically active low and high molecular chemical compounds such as haptens, antigens, antibodies and nucleic acids. Nucleic acids such as DNA or RNA are especially preferred.

Samples according to the invention include clinical samples such as blood, serum, oral rinses, urine, cerebral fluid, sputum, stool, biopsy specimens and bone marrow samples. The sample can also be of a type used for environmental analysis, food analysis or molecular biology research, e.g., from bacterial cultures, phage lysates and products of amplification procedures such as the PCR.

The particles according to the invention have an inner core to which the outer glass surface is applied. The core can be a composite material, or it can be a simple iron core. The core can also consist of a crystalline, ceramic or glass-like structure in which iron oxide is embedded.

The procedure described can be used to isolate native or modified biological material. Native biological material is understood to be material, the structure of which was not irreversibly changed compared with the naturally-occurring biological materials. This does not mean that other components of the sample can not be modified, however. If cells are isolated, for example, the medium surrounding the cells can be modified, but not the cells themselves. If nucleic acids are isolated, they should be cut or modified in their native form, i.e., non-denatured, not cut or not modified by coupling them with reactive groups. The concept of native biological material therefore does not encompass biotinylated nucleic acids in particular. Examples of native biological materials are phage DNA or cellular nucleic acids from blood.

Modified biological materials include materials that do not occur in nature, e.g., nucleic acids that are modified by attaching to them groups that are reactive, detectable or capable of immobilization. An example of this are biotinylated nucleic acids.

In certain cases the sample can be used without pretreatment in the isolation procedure according to the invention. In many cases, however, the sample should be lysed using an appropriate method, releasing the biological material contained in the sample. Procedures for lysing samples are known by the expert and can be chemical, enzymatic or physical in nature. A combination of these procedures is applicable as well. For instance, lysis can be performed using ultrasound, high pressure, by shear forces, using alkali, detergents or chaotropic saline solutions, or by means of proteinases or lipases.

With regard for the lysis procedure to obtain nucleic acids, special reference is made to Sambrook et al.: Molecular Cloning, A Laboratory Manual, 2nd Addition, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, NY and Ausubel et al.: Current Protocols in Molecular Biology 1987, J. Wiley and Sons, NY.

In addition to the biological material to be isolated, the sample can also contain other components in a fluid such as cell residue, proteins, salts and other substances that are not to be isolated. This sample, which preferably contains the biological material in native form, is brought in contact with the particles under conditions in which the target biological material binds to the particle surface. The conditions for this depend on the type of biological material involved, but are basically known. They also depend on the method by which the biological material is bound to the surface. If immunological interactions are utilized for the binding, for instance, conditions must be selected that are suitable for the formation of immunocomplexes. If modified nucleic acids are used, the binding can take place via the groups of nucleic acids that represent the modification, e.g., biotin via binding with streptavidin-coated surfaces. With nucleic acids in particular, however, a direct binding of nucleic acids to glass is preferred because among other reasons the nucleic acids do not have to be modified and even native nucleic acids can be bound. The procedure for binding native nucleic acids to glass particles can be analogous to the procedure described in the prior art. It is preferably performed in the presence of chaotropic salts with a concentration of between 2 and 8 mol/l, and preferably between 4 and 6 mol/l. Chaotropic salts can be sodium iodite, sodium perchlorate, guanidinium thiocyanate, guanidinium isothiocyanate or guanidinium hydrochlorite. Other compounds are also possible.

To bring the sample in contact with the particles, the sample is mixed with the particles and incubated for a period of time sufficient for the binding to occur. Experts are usually familiar with the duration of the incubation step from procedures for performing treatment with non-magnetic particles. This step can be optimized by determining the quantity of immobilized biological material on the surface at different points in time. Incubation times of between 10 seconds and 30 minutes can be appropriate for nucleic acids.

Depending on the size and type of magnetic particles, the particles either separate out of the fluid during the incubation period itself or the suspension remains intact for a longer period of time. If the particles are very small and superparamagnetic, the suspension remains intact for a longer period of time. If the particles are of larger size, the particles slowly separate out of the fluid during the incubation period. Aggregates of this nature form in particular when ferromagnetic particles are involved. When the ferromagnetic particles are not premagnetized, as is preferred, a very gentle separation is guaranteed.

Immobilization is preferably not performed via precipitation by lowering the solubility of the materials to be immobilized. Rather, immobilization is based on biospecific interactions (capture molecules) or adsorption. This largely prevents contaminants from being non-specifically included.

After incubation, the biological material is separated from the fluid. This is achieved in general by separating the material bound to the magnetic particles using a magnetic field. For instance, the magnetic particles can be pulled to the wall of the vessel in which incubation was performed. The fluid containing the sample contents that were not bound to the magnetic particles can then be removed. The removal procedure used depends on the type of vessel in which incubation was performed. Suitable steps include removing the fluid via pipetting or aspiration.

The magnetic particles can then be purified one or more times using a wash solution, if desired. A wash solution is used that does not cause the biological material to be deliberated from the particle surface but that washes away the undesired contaminants as thoroughly as possible. This wash step preferably takes place by incubating the wash solution with the particles. The particles are preferable resuspended during this step, e.g., by means of shaking or applying a magnetic field that is not identical to the first magnetic field. The contaminated wash solution is preferably separated just like the sample in the step described above for binding the biological material.

After the last wash step, the magnetic particles can be dried briefly in a vacuum, or the fluid can be allowed to evaporate. A pretreatment step using acetone may also be performed.

If desired, the biological material purified in this manner can be separated from the magnetic particles. This step also depends on the manner in which the biological material was bound to the magnetic particles. If the biological material is native nucleic acids and the magnetic particles are glass-coated particles, the nucleic acids can be removed from the particles according to the invention using an elution buffer having a low salt content. Buffers of this nature are known from DE 3724442 and Analytical Biochemistry 175, 196-201 (1988). The elution buffers with a low salt content are in particular buffers with a content of less than 0.2 mol/l. In an especially preferred embodiment, the elution buffer contains Tris. In another special embodiment, the elution buffer is demineralized water.

In yet another embodiment, the purification and isolation procedure described is performed after the cells (e.g., viral particles or prokaryotic or eukaryotic cells) are separated immunomagnetically from a bodily fluid or tissue. In this step, the sample is incubated, e.g., while shaking, with magnetic particles to which an antibody against an antigen on the cell is immobilized. These particles can be particles according to the invention or commercially available particles (e.g., MACS Microbeads from Miltenyi Biotec GmbH, Bergisch Gladbach, Germany). After a magnetic field is applied, one or more wash steps are performed using a saline solution. Particles are obtained to which the desired cells are bound. The bound cells are then resuspended in a saline buffer. In a preferred embodiment, this saline buffer is a chaotropic saline solution so that the nucleic acids contained in the cell are released from the cells.

An especially advantageous procedure for isolating nucleic acids from samples containing cells is achieved by combining the isolation of cells described above with the isolation of nucleic acids — preferable in their native form — also described above, on the magnetic particles according to the invention. The advantage of this embodiment is

its potential simplicity (single-tube method), high sensitivity (especially important in medical microbiology and oncology), and the ease with which it can be automated.

The biological materials isolated using the procedure according to the invention can now be used further as necessary. For instance, they can be used as a substrate for various enzymatic reactions. When nucleic acids are involved, they can be used for sequencing, radioactive or non-radioactive labelling, amplification of one or more of the sequences they contain, transcription, hybridization with labelled probe nucleic acids, translation or ligation. An advantage of the procedure according to the invention is that it is very easy to separate the biological material from the fluid. In the prior art, a centrifugation step was used to separate the glass particles from contaminants, or, when the biological material is bound to glass fiber filters, the fluid is drawn through the filters. This is a limiting step that makes it difficult to process large quantities of sample.

The biological materials can be separated from contaminants more effectively using the particles according to the invention. In particular, inhibitors for certain enzymatic reactions can be removed to a large extent according to the invention. The yield of biological material is relatively high. Fractionation of long nucleic acids was not observed. The particles according to the invention can preferably be magnetized more quickly.

Figure 1 illustrates the isolation of nucleic acids from a sample containing cells.

Figure 2 illustrates the separation of isolated nucleic acids according to the invention in an agarose gel.

Figure 3 depicts the separation of reaction products after isolation according to the invention and amplification by means of the PCR.

Figure 4 shows a gel with the results from example 4.

Figure 1 illustrates the isolation of nucleic acids from a sample containing cells. The sample (specimen) that contains cells is pretreated in a sample-specific fashion so that the cells in which the nucleic acids are to be detected are present in the proper form.

When samples are used from which bodily fluids were removed, for instance, this entails adding reagents, e.g., to liquify viscous samples such as saliva. An antibody bound to a solid phase, preferably a bead, that can detect and bind the cell is added to a vessel containing the sample treated in this fashion. Antigens on the cell surface have proven to be suitable partners for the antibody, for instance. The specificity of the antibody can depend on the specificity of the analysis to be performed. If the solid phase is the wall of the vessel, the cells are bound directly to the wall. If the solid phase is comprised of beads, they are separated from the fluid using suitable separation methods. This can be performed by means of filtration, for instance. If magnetic beads are used, they can be separated out by applying a magnetic field to the outside wall of the vessel. The separated cells are washed with a fluid to remove contaminants (that would interfere with the detection) along with the medium surrounding the cells. The conditions are preferably such that the cells are neither separated from the solid phase nor destroyed. The cells are then destroyed, i.e., lysed. This can be performed, for instance, by treating the cells with chaotropic salts. Other possibilities include the application of proteinases and detergents.

In the preferred embodiment, the particles according to the invention are added to the lysis mixture. After a suitable period of time for the lysis to take place — which can be optimized by loading the surface with nucleic acids — the particles are separated from the surrounding fluid that contains additional cell components that are not to be detected. This is performed preferably by applying a magnetic field by placing a magnet against the vessel wall.

To remove any contaminants that may still be present, a wash step is preferably performed with a fluid that does not cause the nucleic acids to be determined to be separated from the glass surface. An elution buffer having reagent conditions under which the nucleic acids separate from the glass surface is added to remove the nucleic acids from the glass surface. These conditions are low salt conditions in particular. Depending on the intended further use of the nucleic acids, the fluid can now be separated from the particles and processed further. This separation step is preferably

performed via application of a magnetic field so that the particles are separated from each other.

The following examples explain the invention in greater detail.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210

Example 1Manufacture of the magnetic particles according to the invention

Six different sols were used. The sols were manufactured as follows:

Sol 1 ($\text{SiO}_2:\text{B}_2\text{O}_3 = 7:3$):

Synthesis was performed in a 250 ml round flask while stirring constantly.

86.6 ml tetraethyl orthosilicate

+ 7 ml anhydrous, non-denatured ethanol

+ 14.1 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

+ 37.8 ml trimethylborate

then keep the sol at 50°C for 2 hours. Add

+ 14.1 ml 0.15 M HCl

Sol 2 ($\text{SiO}_2:\text{B}_2\text{O}_3 = 4:1$):

Synthesis was performed in a 250 ml round flask while stirring constantly.

100.5 ml tetraethyl orthosilicate

+ 7 ml anhydrous, non-denatured ethanol

+ 16.3 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

+ 25.6 ml trimethylborate

then keep the sol at 50°C for 2 hours. Add

+ 16.3 ml 0.15 M HCl

Sol 3 ($\text{SiO}_2:\text{B}_2\text{O}_3 = 85:15$):

Synthesis was performed in a 250 ml round flask while stirring constantly.

107.8 ml tetraethyl orthosilicate

- 16 -

- + 7 ml anhydrous, non-denatured ethanol
- + 17.5 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

- + 19.4 ml trimethylborate
- then keep the sol at 50°C for 2 hours. Add
- + 17.5 ml 0.15 M HCl

Sol 4 ($\text{SiO}_2:\text{B}_2\text{O}_3 = 4:1$; 2 Mol% P_2O_5):

Synthesis was performed in a 250 ml round flask while stirring constantly.

- 100.5 ml tetraethyl orthosilicate
- + 7 ml anhydrous, non-denatured ethanol
- + 16.3 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

- + 25.6 ml trimethylborate
- then keep the sol at 50°C for 2 hours. Add
- + 16.3 ml 0.15 M HCl
 - + 1.63 g P_2O_5

Sol 5 ($\text{SiO}_2:\text{B}_2\text{O}_3 = 4:1$ Mol% Al_2O_3):

Synthesis was performed in a 250 ml round flask while stirring constantly.

- 100.5 ml tetraethyl orthosilicate
- + 7 ml anhydrous, non-denatured ethanol
- + 16.3 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

- + 25.6 ml trimethylborate
- then keep the sol at 50°C for 2 hours. Add

+ 16.3 ml 0.15 M HCl

+ 3.06 g AlCl_3

Sol 6 ($\text{SiO}_2\text{:B}_2\text{O}_3 = 4:1$ Mol% ZrO_2):

Synthesis was performed in a 250 ml round flask while stirring constantly.

100.5 ml tetraethyl orthosilicate

+ 7 ml anhydrous, non-denatured ethanol

+ 16.3 ml 0.15 M HCl

A biphasal mixture is produced. Stir it at room temperature until it becomes a single phase. Add dropwise

+ 25.6 ml trimethylborate

+ 5.15 ml zircon(IV)-proylate, 70% solution by weight in 1-propanol

then keep the sol at 50°C for 2 hours. Add

+ 16.3 ml 0.15 M HCl

After another 2 hours at 50°C, 22.5 g Iriodin 600 (black mica) was added for each 150 ml sol and stirred. It was then coated with a spray dryer (Büchi 190, Mini Spray Dryer). The temperature of the spray dryer nozzle was 134°C.

The powder obtained in the spray drying process was then subjected to a temperature treatment step in a nitrogen atmosphere (90 l/h). The temperature was increased at a rate of 1 k/min and the powder was maintained at a densification temperature for 2 hours. For coating with sol 1, this temperature was 750°C, and 860°C for coating with sol 2. The temperature was 800°C for all other coating processes. After the sintering process the oven was turned off and the powder was brought to room temperature. Agglomerates were sifted out using a 50 μm sieve.

Example 2Manufacture of GMP1, GMP2, GMP3 and GMP4

GMP1, GMP2, GMP3 and GMP4 are pigments from different production lots that were obtained from sol 1 (example 1) in a process described in example 1, under the following conditions:

| Parameter | GMP 1 | GMP 2 | GMP 3 | GMP 4 |
|---|---------|---------|---------|---------|
| Aging of the sol (h) (30°C) | 36 | 36 | 36 | 36 |
| Percentage of pigment in sol (g/100 ml) | 5 | 15 | 8 | 20 |
| Nozzle air flow (%) | 100 | 100 | 100 | 100 |
| Air pressure (bar) | 6 | 6 | 6 | 3 |
| Nozzle temperature (°C) | 135 | 120 | 130 | 143 |
| Densification temperature (°C) | 534 | 534 | 534 | 615 |
| subsequent O ₂ -treatment (1 hour) | (300°C) | (300°C) | (300°C) | (400°C) |
| Pigment yield | low | high | medium | high |
| DNA yield | low | high | high | high |

Example 3PCR sample pretreatment from human whole blood using magnetic glass particles**Nucleic Acid Isolation**

10 mg each from 3 lots of glass magnetic particles (GMP 2-4) were placed in Eppendorf test tubes. The exact sample weights are indicated in Table 1. Three-fold determinations were performed.

40 µl proteinase K (20 mg/ml, made from lyophilisate) were added via pipetting to each 200 µl of thawed whole blood and mixed immediately. In the next step, 200 µl binding buffer (6 M guanidine-HCl, 10 mM Tris-HCl, 10 mM urea, 30% Triton X-100, pH 4.4) were added, mixed, and then incubated for 10 minutes at 70°C. 200 µl i-propanol were added, and the preparation was then mixed on the vortex mixer for 10 seconds. The sample was left at room temperature for 20 minutes, then mixed once more for 10 seconds. The magnetic separation step was performed for at least 30 seconds in a

magnetic particle separator from Boehringer Mannheim (ID# 1 641 794). The supernatant was removed and analyzed as described below.

The magnetic particles were washed with 500 μ l wash buffer (20 mM NaCl, 10 mM Tris-HCl, pH 7.5 (25°C), 80% ethanol) by mixing for 10 seconds, leaving them at room temperature for 1 minute, then mixing for 10 seconds. They were then pulled to the vessel wall using the magnetic particle separator. The supernatant was removed and discarded. The wash procedure was repeated until the wash fluid was colorless (4 times in all). The nucleic acids were then eluted 3 x with 200 μ l each time of elution buffer prewarmed to 70°C (10 mM Tris-HCl, pH 8.5), then mixed for 10 seconds, left at room temperature for 10 minutes, and mixed for 10 minutes.

Preparing the Supernatant

The supernatant obtained after the first binding to the magnetic glass particles was investigated as follows for its nucleic acid content: the supernatant was placed in a filter tube (Boehringer Mannheim ID# 1744003, as provided in the *High Pure* PCR Product Purification Kit, for instance) and centrifuged for 1 hour at 8000 rpm in an Eppendorf tabletop centrifuge. The flow-through material was discarded and the filter tube was washed 2 x with 500 μ l wash buffer (centrifugation as described above). The filter tube was centrifuged briefly to dryness, and then eluted with 2 x 200 μ l 1 x elution buffer prewarmed to 70°C by centrifuging once more.

Analyzing the Eluate and Sample Supernatant

10 μ l of sample buffer were added to 50 μ l of the eluate and the supernatant prepared using the filter tube, respectively. 45 μ l of this preparation were separated in an 0.8% agarose gel using electrophoresis at 120 V for 90 minutes.

Various dilutions of the eluate and the prepared supernatants were measured using spectroscopy at 260 and 280 nm in a Uvikon 710 (Kontron).

Two 5 μ l aliquots of eluate were investigated in duplicate determinations using Expand™ Long Template PCR (Boehringer Mannheim ID# 1681834) with specific primers for the human tPA gene (expected length of product: 15 kb).

| Mix I | per batch | Mix II | per batch |
|----------------------|------------------|----------------------|------------------|
| dNTP, 100 mM each | 1 μ l | Expand™ buffer, 10 x | 5 μ l |
| Primer 1, 200 ng/ ml | 1 μ l | Expand™ polymerase | 0.75 μ l |
| Primer 2, 225 ng/ ml | 1 μ l | H2O, bidistilled | 19.25 μ l |
| H2O, bidistilled | 17 μ l | | |
| | <hr/> 20 μ l | | <hr/> 25 μ l |

Mix I is placed in a thin-walled PCR tube with 5 μ l eluate, then mix II is added. The preparation is mixed briefly, then covered with a layer of 30 μ l of mineral oil. The preparations are amplified in a Perkin Elmer thermal cycler 9600 with the following settings:

| | | |
|----------------------|------|-----------|
| 2 minutes | 92°C | |
| 10 seconds | 92°C | |
| 30 seconds | 65°C | 10 cycles |
| 12 minutes | 68°C | |
| 10 seconds | 92°C | |
| 30 seconds | 65°C | 20 cycles |
| 12 minutes + | 68°C | |
| 20 seconds per cycle | | |
| 7 minutes | 68°C | |

then

7°C

10 µl sample buffer were added to the 50 µl PCR preparations. 45 µl of this mixture were then separated in an 0.8% agarose gel using electrophoresis at 120 V for 90 minutes.

Results

| | | | Super natant | | 1:8 | 1. Eluate | | 1:8 | | |
|-------|-------|---|-----------------|--------|--------|--------------|-----------|-----------|---------|---------|
| | | | 260 nm | 280 nm | Yield | 260/280 | 260 nm | 280 nm | Yield | 260/280 |
| | | | | | | 0 | | | | 0 |
| GMP/2 | 12 mg | 1 | 0,021 | 0,013 | 1,7 µg | 1,6 | 0,171 | 0,164 | 13,7 µg | 1,0 |
| | 10 mg | 2 | 0,045 | 0,035 | 3,7 µg | 1,3 | 0,137 | 0,138 | 11,0 µg | 1,0 |
| | 9 mg | 3 | 0,036 | 0,027 | 2,9 µg | 1,3 | 0,153 | 0,164 | 12,2 µg | 0,9 |
| GMP/3 | 10 mg | 1 | 0,050 | 0,042 | 4,0 µg | 1,2 | 0,245 | 0,246 | 19,6 µg | 0,9 |
| | 10 mg | 2 | 0,033 | 0,022 | 2,6 µg | 1,5 | 0,397 | 0,398 | 31,8 µg | 1,0 |
| | 10 mg | 3 | 0,042 | 0,030 | 3,4 µg | 1,4 | 0,278 | 0,282 | 22,2 µg | 0,9 |
| GMP/4 | 10 mg | 1 | 0,065 | 0,056 | 0,7 µg | 1,2 | 0,135 | 0,142 | 11,0 µg | 1,0 |
| | 11 mg | 2 | 0,071 | 0,142 | 2,4 µg | 0,5 | 0,140 | 0,142 | 11,2 µg | 1,0 |
| | 10 mg | 3 | 0,066 | 0,051 | 1,7 µg | 1,3 | 0,130 | 0,130 | 10,4 µg | 1,0 |

| | | | 2. Eluate | | 1:8 | 3. Eluate | | 1:4 | | |
|-------|---|-------|-----------|---------|-------|--------------|-----------|-----------|-------|-------------|
| | | | 260 nm | 280 nm | Yield | 260/280 | 260 nm | 280 nm | Yield | 260/280 |
| | | | | | | | | | | Σ Eluate |
| GMP/2 | 1 | 0,099 | 0,101 | 7,9 µg | 1,0 | 0,057 | 0,062 | 2,3 µg | 0,9 | 23,9 µg |
| | 2 | 0,078 | 0,076 | 6,2 µg | 1,0 | 0,041 | 0,049 | 1,6 µg | 0,8 | 18,8 µg |
| | 3 | 0,103 | 0,112 | 8,2 µg | 0,9 | miss | | | | |
| GMP/3 | 1 | 0,147 | 0,147 | 11,8 µg | 1,0 | 0,084 | 0,098 | 3,4 µg | 0,9 | 34,8 µg |
| | 2 | 0,256 | 0,252 | 20,5 µg | 1,0 | 0,042 | 0,043 | 1,7 µg | 1,0 | 54,0 µg |
| | 3 | 0,147 | 0,143 | 11,8 µg | 1,0 | 0,073 | 0,093 | 2,9 µg | 0,8 | 36,9 µg |
| GMP/4 | 1 | 0,106 | 0,108 | 8,5 µg | 1,0 | 0,083 | 0,098 | 3,3 µg | 0,8 | 22,8 µg |

| | | | | | | | | | |
|---|-------|-------|---------|-----|-------|-------|--------|-----|---------|
| 2 | 0,111 | 0,114 | 8,9 µg | 1,0 | 0,054 | 0,063 | 2,2 µg | 0,9 | 22,3 µg |
| 3 | 0,135 | 0,141 | 10,8 µg | 1,0 | 0,077 | 0,095 | 3,1 µg | 0,8 | 24,3 µg |

Table 1: Yield of nucleic acids using magnetic glass particles and 200 µl blood

The first eluates were still slightly yellow in color and slightly contaminated with fine magnetic particles.

The analysis of the eluates in agarose gel (Fig. 2) reveals good reproducibility of the yield. The magnetic particles GMP 2-4 show no significant differences. Eluates 1 (above) and 2 (below) contain approximately the same concentration of nucleic acids (estimated by the gel). Eluate 3 has a low concentration of nucleic acids. The supernatants also contain a low concentration of nucleic acids.

The Expand™PCR yields very good, specific amplification products for all samples, with just a few outliers (Table 2). When magnetic glass beads are used, nucleic acids are isolated from human blood samples that then yielded specific amplicates in a subsequent PCR step.

| | | 15 kb Expand™PCR | | Human tPA Gene | |
|-------------------|---|------------------|-----|----------------|------|
| | | 1st Eluate | | 2nd Eluate | |
| GMP/2 | 1 | | n/a | + | + |
| | 2 | + | + | + | + |
| | 3 | + | + | | n/a |
| GMP/3 | 1 | + | + | + | + |
| | 2 | (+) | + | + | + |
| | 3 | - | (+) | + | + |
| GMP/4 | 1 | + | + | + | + |
| | 2 | + | + | + | (+)* |
| | 3 | + | + | | n/a |
| K, BM Control DNA | | | | | |

Table 2. Results with Expand™PCR

*3rd eluate

Fig. 3 shows a gel with the reaction products after PCR amplification. MWV III is a molecular weight marker (eluate 1, above; eluate 2, below).

Example 4

Binding of DNA length standard to magnetic glass particles

1. Preparation of the magnetic glass particles

12 mg of glass magnetic particles from GMP 4 are placed in a 12 mg Eppendorf test tube.

2. Lysis and binding

900 µl lysis buffer (4.6 M GuSCN, 45 mM Tris, 20 EDTA, pH 7.3) and 100 µl DNA sample in which DNA length standard III from Boehringer Mannheim (Cat. No. 528552) was added as a model are mixed in a 1.5 ml Eppendorf vessel with 12 mg magnetic glass particles for 2 to 10 seconds until a homogenous suspension is obtained. The solution is incubated at room temperature for 20 minutes and mixed every 5 minutes.

Magnetic separation is performed for at least 15 seconds in a magnetic particle separator. The supernatant is removed via pipetting.

3. Washing and Drying

The magnetic glass particles are washed twice with wash buffer (5.2 M GuSCN, 50 mM Tris, pH 6.5), twice with 70% precooled ethanol, and once with acetone by removing the magnetic field, adding 800 μ l solution via pipetting, mixing for 2 seconds, leaving at RT for 1 minutes, applying the magnetic field and then removing the supernatant via pipetting.

When the acetone is removed, the particles are dried for 10 minutes at 56°C in the heating block with the cover open.

4. Eluting the DNA

The DNA is eluted with 4 x 50 μ l elution buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) by incubating it at 56°C for 10 minutes while shaking repeatedly. The supernatant, which contains the DNA, is then transferred to a new Eppendorf vessel via pipette.

5. Analyzing the Eluate

Sample buffer is added to one-fifth of the eluate volume and the DNA is separated on a 1% agarose gel at 90 V. To determine the recovery, a dilution series of DNA length standard III is applied to the same gel that contains the quantities of DNA expected in the sample.

The quantitative evaluation is performed by scanning a Polaroid photo of the agarose gel. The dilution series of the standard is used as the calibrator.

The yield of DNA using magnetic glass particles is shown in Table 1.

| Standard DNA - amount | | | Intensity. | Sample | Type of pigment/ bead | Intensity sample (measured) [rel.Units] | calculated amount of DNA- in the gel [ng] | calculated amount of DNA in the sample [ng] | Recovery [%] |
|-----------------------|-----------------|-----------------------------|------------------------------------|--------|--------------------------|---|---|---|-----------------|
| No. | in the standard | Standard (measured) [ng] | Standard (measured) [rel.Units] | | | | | | |
| 1 | 200 | 65 | 1 | GMP4 | 45 | 139 | 695 | 69,5 | |
| 2 | 175 | 56 | 2 | GMP4 | 39 | 120 | 600 | 60,0 | |
| 3 | 150 | 51 | | | | | | | |
| 4 | 125 | 44 | | | | | | | |
| 5 | 100 | 37 | | | | | | | |
| 6 | 75 | 25 | | | | | | | |
| 7 | 50 | 17 | | | | | | | |
| 8 | 25 | 9 | | | | | | | |
| 9 | 10 | 4 | | | | | | | |

Table 1. Yield of DNA length standard III with magnetic glass particles

The agarose gel that was used as the basis for the quantitative evaluation is shown in Fig. 4. It is a 1% ethidium bromide-stained agarose gel. Lanes 1 through 10 correspond to a dilution series of DNA length standard III. 1: 1 µg DNA, 2: 200 ng DNA, 3: 175 ng DNA, 4: 150 ng DNA, 5: 125 ng DNA, 6: 100 ng DNA, 7: 75 ng DNA, 8: 50 ng DNA, 9: 25 ng DNA, 10: 10 DNA.

Lanes 11 and 12 correspond to the DNA eluted from the magnetic glass particles with 200 ng DNA length standard added.

SEQUENCE PROTOCOL

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Boehringer Mannheim GmbH

(B) STREET: Sandhoferstr. 116

(C) CITY: Mannheim

(D) COUNTRY: DE

(E) POSTAL CODE: 68298

(F) TEL.: 0621 759 4348

(G) FAX: 0621 759 4457

(ii) NAME OF THE INVENTION: Magnetic Pigment

(iii) NUMBER OF SEQUENCES: 2

(iv) COMPUTER-READABLE FORM:

(A) DATA CARRIER: Floppy disk

(B) COMPUTER: IBM PC Compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPA)

(2) INFORMATION ON SEQ ID# 1:

(i) SEQUENCE IDENTIFICATION:

(A) LENGTH: 34 base pairs

(B) TYPE: Nucleotide

(C) STRAND TYPE: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: /desc = "oligodeoxyribonucleotide"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ACTGTGCTTC TTGACCCATG GCAGAAGCGC CTTC 34

(2) INFORMATION ON SEQ ID NO: 2:

(i) SEQUENCE IDENTIFICATION: (A) LENGTH: 34 base pairs

(B) TYPE: Nucleotide

(C) STRAND TYPE: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: /desc = "oligodeoxyribonucleotide"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CCTTCACTGT CTGCCTAACT CCTTCGTGTG TTCC 34

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000